


U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER BARENHOLZ=6
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NO. (if known, see 37 CFR 1.5) 10/009771
INTERNATIONAL APPLICATION NO. PCT/IL00/00350	INTERNATIONAL FILING DATE 15 June 2000	PRIORITY CLAIMED 15 June 1999
TITLE OF INVENTION ENZYMATIC PREPARATION OF PHOSPHOLIPIDS IN AQUEOUS MEDIA		
APPLICANT(S) FOR DO/EO/US Yechezkel BARENHOLZ et al.		
<p>Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:</p> <ol style="list-style-type: none"> 1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). 4. <input checked="" type="checkbox"/> The US has been elected in a Demand by the expiration of 19 months from the priority date (PCT Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> a. <input type="checkbox"/> is attached hereto (required only if not transmitted by the International Bureau). b. <input checked="" type="checkbox"/> has been communicated by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). <p>Items 11. to 16. below concern document(s) or information included:</p> <ol style="list-style-type: none"> 11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input type="checkbox"/> An Assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> A substitute specification. 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input checked="" type="checkbox"/> Other items or information: <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Courtesy copy of the International Application as filed. <input checked="" type="checkbox"/> Courtesy copy of the first page of the International Publication (WO 00/77183). <input checked="" type="checkbox"/> Courtesy copy of the International Preliminary Examination Report. There were no annexes <input checked="" type="checkbox"/> Courtesy Copy of the International Search Report. <input checked="" type="checkbox"/> Application Data Sheet. <p><input checked="" type="checkbox"/> The application is (or will be) assigned to: YISSUM RESEARCH DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEM, whose address is Jabotinsky Street 46, 91042 Jerusalem, Israel.</p>		

JC07 Rec'd PCT/PTO 1 7 DEC 2001

U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 10/009771		International Application No. PCT/IL00/00350		Attorney's Docket No. BARENHOLZ=6	
17. [xx] The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a)(1)-(5): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00				CALCULATIONS PTO USE ONLY	
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$ 890.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than [] 20 [X] 30 months from the earliest claimed priority date (37 CFR 1.492(c)).				\$ 130.00	
Claims as Originally Presented	Number Filed	Number Extra	Rate		
Total Claims	14 - 20		X \$18.00	\$	
Independent Claims	1 - 3		X \$84.00	\$	
Multiple Dependent Claims (if applicable)			+ \$280.00	\$	
TOTAL OF ABOVE CALCULATIONS =				\$ 990.00	
Claims After Post Filing Prel. Amend	Number Filed	Number Extra	Rate		
Total Claims	- 20		X \$18.00	\$	
Independent Claims	- 3		X \$84.00	\$	
TOTAL OF ABOVE CALCULATIONS =				\$ 990.00	
Reduction of ½ for filing by small entity, if applicable. Applicant claims small entity status. See 37 CFR 1.27.				\$ 495.00	
SUBTOTAL =				\$ 495.00	
Processing fee of \$130.00 for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =				\$ 495.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$	
TOTAL FEES ENCLOSED =				\$ 495.00	
				Amount to be:	\$
				refunded	\$
				charged	\$
a. [] A check in the amount of \$ _____ to cover the above fees is enclosed. b. [X] Credit Card Payment Form (PTO-2038), authorizing payment in the amount of \$ 495.00, is attached. c. [] Please charge my Deposit Account No. 02-4035 in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed. d. [XX] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 02-4035 . A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO:					
BROWDY AND NEIMARK, P.L.L.C. 624 NINTH STREET, N.W., SUITE 300 WASHINGTON, D.C. 20001 TEL: (202) 628-5197 FAX: (202) 737-3528 Date of this submission: Monday, December 17, 2001					
Form PTO-1350 (as slightly revised by Browdy and Neimark)					


 SIGNATURE
 Roger L. Browdy
 NAME
 25,618
 REGISTRATION NUMBER

INVENTOR INFORMATION

JCN 7 1000000771 121302
Rec'd PCT/PTO 17 DEC 2001
10/009771

Inventor One Given Name:: Yechezkel
Family Name:: BARENHOLZ
Postal Address Line One:: Nave Shanan Street 18
City:: Jerusalem
Country:: Israel
Postal or Zip Code:: 93707
Citizenship Country:: Israel
Inventor Two Given Name:: Shimon
Family Name:: AMSELEM
Postal Address Line One:: Miriam Mizrachi Street 10
City:: Rehovot
Country:: Israel
Postal or Zip Code:: 76551
Citizenship Country:: Israel

CORRESPONDENCE INFORMATION

Correspondence Customer Number:: 001444
Fax One:: 202-737-3528
Electronic Mail One:: Mail@BrowdyNeimark.com

APPLICATION INFORMATION

Title Line One:: ENZYMATIC PREPARATION OF PHOSPHOLIPIDS I
Title Line Two:: N AQUEOUS MEDIA
Formal Drawings?: No
Docket Number:: BARENHOLZ-6
Secrecy Order in Parent Appl.?: No

REPRESENTATIVE INFORMATION

Representative Customer Number:: 1444

CONTINUITY INFORMATION

This application is a:: 371 OF
> Application One:: PCT/IL00/00350
Filing Date:: 06-15-2000

Which is a:: NON PROV. OF PROVISIONAL
>> Application Two:: 60/139316
Filing Date:: 06-15-1999

Source:: PrintEFS Version 1.0.1

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)	Art Unit:
Yechezkel BARENHOLZ et al.)	
)	
IA No.: PCT/IL00/00350)	
)	Washington, D.C.
IA Filed: June 15, 2000)	
)	
U.S. App. No.:)	
(Not Yet Assigned))	
)	December 17, 2001
National Filing Date:)	
(Not Yet Received))	
)	
For: ENZYMATIC PREPARATION..)	Docket No.:
		BARENHOLZ=6

PRELIMINARY AMENDMENT

Honorable Commissioner for Patents and Trademarks
Washington, D.C. 20231

Sir:

Contemporaneous with the filing of this case and
prior to calculation of the filing fee, kindly amend as
follows:

IN THE SPECIFICATION

After the title please insert the following
paragraph:

REFERENCE TO RELATED APPLICATIONS

The present application is the national stage under
35 U.S.C. 371 of international application PCT/IL00/00350,
filed June 15, 2000 which designated the United States, and

In re of: Yechezkel BARENHOLZ et al. (BARENHOLZ=6)


which international application was published under PCT
Article 21(2) in the English language.

REMARKS

The above amendment to the specification is being
made to insert reference to the PCT application of which the
present case is a U.S. national stage.

Favorable consideration is earnestly solicited.

Respectfully submitted,
BROWDY AND NEIMARK, P.L.L.C.
Attorneys for Applicant

By: 
Roger L. Browdy
Registration No. 25,618

RLB:wrđ
Telephone No.: (202) 628-5197
Facsimile No.: (202) 737-3528
F:\,C\ohn\Barenholz6\PTO\Preliminary Amendment.doc

Field of the Invention

The present invention relates to method of conducting a phospholipase-catalyzed transesterification or hydrolysis of a phospholipid. The reaction is useful for the preparation of a variety of differently substituted phospholipids, as well as diacyl glycerols and ceramides. The method employs a water/solid particle interface, gives high conversions, and does not require the use of organic solvents, detergents or surfactants.

References

Allgyer, T.T. and Wells, M.A., *Biochemistry* **18**:5348 (1979).

Amselem, S. *et al.*, *Chem. Phys. Lipids* **64**, 219-237 (1993).

Barenholz, Y. and Amselem, S., "Liposome Preparation and Related Techniques", pp. 527-616, in *Liposome Technology*, 2nd ed., Vol. 1, G. Gregoriadis, ed., CRC Press, Boca Raton, FL, 1993.

Bartlett, G.R., *J. Biol. Chem.* **234**:466-71 (1959).


Bligh, E.G. and Dyer, W.J., *Can. J. Biochem.* **31**:911 (1959).

Servi, S., *Topics in Current Chemistry* **200**:127-158 (1999).

Yang, S.F., *Methods in Enzymol.* **14**:208-211 (1969).

Background of the Invention

Phospholipases A1, A2, C, and D enzymatically cleave the bonds in a diacyl phospholipid as illustrated below:



These enzymes have proven useful in the synthetic preparation of a wide variety of phospholipids, sphingolipids, and neutral lipids, and in particular the preparation of new, non-naturally occurring, or non-abundant lipids from more abundant source materials. Similarly, C- and D-type phospholipases which use sphingomyelin as a substrate (sphingomyelin phosphodiesterases), and various glycosidases which use glycolipids as substrates, can be utilized to prepare ceramides and other lipids.

In particular, phospholipase D has been used to convert phosphatidyl cholines (PC) to less common lipids such as phosphatidyl serines (PS) and phosphatidyl glycerols (PG). The reaction is a

transesterification (also referred to as transphosphatidylolation) between the starting material, PC, and a hydroxyl containing reagent such as glycerol or serine. The enzyme is especially useful in this respect in that it is able to produce the transesterified product in the presence of more than a stoichiometric amount of water (Servi). An organic solvent or detergent is typically required, as the lipid substrate is not water-soluble, and when lipid is presented to the enzyme in the form of an aqueous liposomal dispersion, the enzymatic activity is usually poor. Conventional methods for this reaction have generally employed a two-phase system containing water, in which the enzyme is soluble, and an organic solvent, in which the lipid is soluble, with the enzymatic reaction occurring at the interface between the water and the organic solvent.

The use of detergents or organic solvents is problematic in large scale reactions, and may be prohibited in the production of many pharmaceutical and food products. In addition, some amount of the hydrolysis product, phosphatidic acid (PA), is typically formed in such reactions, and methods are sought for increasing conversion to the transesterification product.

Summary of the Invention

The present invention includes, in one aspect, a method of conducting an enzyme-catalyzed transesterification or hydrolysis of a phospholipid or glycolipid. The method comprises dissolving the enzyme in an aqueous medium containing a liposomal suspension of the lipid and a hydroxyl-containing reagent, adding silica gel to the medium, and agitating the resulting mixture. For some enzymes, a divalent metal cation, such as Ca^{+2} , Zn^{+2} , or Mg^{+2} , is required for reaction. The enzyme is preferably a phospholipase, such as phospholipase A1, A2, C, or D, or a sphingomyelin phosphodiesterase. In a preferred embodiment, the phospholipase is phospholipase D.

The method is particularly useful for reactions of phospholipids which make up naturally occurring phospholipid mixtures, such as phosphatidyl choline, or for phospholipids isolated from tissue, e.g. a brain phospholipid extract. The hydroxyl-containing reagent can be water, in the case of a hydrolysis reaction; in preferred embodiments, the reagent is an alcohol or alcohol derivative, such as glycerol, serine, inositol, or hydroxy-terminated PEG (polyethylene glycol). Synthetic phospholipids of various stereochemistries can be prepared by varying the stereochemistry of the hydroxyl-containing reagent, e.g. in the case of serine.

In the reaction, the silica gel is preferably added in an amount which is at least four times the amount of the lipid by weight, giving a silica gel/lipid ratio of at least 4:1, and more preferably an amount which is at least ten times the amount of the lipid by weight, giving a silica gel/lipid ratio of at least 10:1. The silica gel preferably has a mean particle size no greater than $25\mu\text{m}$, and more preferably no greater than $15\mu\text{m}$.

In further preferred embodiments, the concentration of phospholipase in the medium is at least

3 mg/ml, and more preferably at least 7 mg/ml. When the enzyme is phospholipase D, the concentration of calcium ion in the medium is preferably in the range of 5-100 mM.

These and other objects and features of the invention will be made more fully apparent in the following detailed description of the invention.

5

Detailed Description of the Invention

I. Definitions

The terms below have the following meanings unless indicated otherwise.

10 A "phospholipid" refers to an amphipathic lipid having one or two hydrophobic acyl chains and a phosphate-containing polar head group. The lipid may contain a chemically reactive group, such as an amine, acid, ester, aldehyde or alcohol, at the polar head group. Synthetic phospholipids may also have a chromophore or a fluorophore attached at various parts of the molecule.

15 The hydrocarbon chains are typically between about 2-26, and preferably about 14-22, carbon atoms in length, and commonly have varying degrees of unsaturation. The hydrocarbon chains may include branching or other modifications, e.g. cyclopropyl or cyclohexyl groups.

Representative examples of phospholipids are phosphatidyl choline (PC), phosphatidyl ethanolamine (PE), phosphatidic acid (PA), phosphatidyl inositol (PI), phosphatidyl glycerol (PG), and sphingomyelin (N-acyl sphingosylphosphocholine, a sphingolipid). The backbone in
20 sphingolipids is derived from sphingosine, rather than from glycerol. In sphingomyelin, the amino group of sphingosine is linked to a fatty acid chain via an amide bond, and the primary hydroxyl group is esterified to phosphoryl choline.

An unusual type of phospholipid is found in the membranes of archaea, a type of bacteria. The cell membranes of these organisms are composed of phytanyl lipids, rather than fatty acids.
25 The phytanyl (3,7,11,15-tetramethylhexadecyl) lipids are joined to glycerol to form lipids by an ether rather than an ester bond.

The name of a particular lipid includes the specific acyl groups, e.g. dimyristoyl phosphatidyl choline (DMPC) or N-palmitoyl sphingomyelin. Naturally occurring lipid mixtures are common, and these include the lecithins, such as soy lecithin or egg lecithin, which are mixtures of
30 phosphatidyl choline having various acyl chains. Such mixtures may be used in the present reaction, as well as partially purified tissue extracts.

A "phospholipase" is an enzyme that cleaves (by hydrolysis or transesterification) ester bonds in phospholipids. The two general types are aliphatic esterases (types A1, A2 and B), which release fatty acids from glycerol-based phospholipids, and phosphodiesterases (types C and D), which
35 cleave phosphate ester bonds. Sphingomyelinases (sphingomyelin phosphodiesterases) may be similar or identical to phospholipases which act on glycerophospholipids, as in many bacterial

enzymes, or they may be specific for sphingolipids, as in eukaryotic enzymes. As used herein, the category of C- and D-type phospholipases includes sphingomyelin phosphodiesterases.

"Silica gel" refers to a colloidal, highly adsorbent form of silicon dioxide or a salt (silicate) thereof. The term as used herein covers commercial silica gels used for chromatography, preferably TLC grade, which may comprises silicic acid ($\text{SiO}_2 \cdot n\text{H}_2\text{O}$) or a salt such as magnesium silicate (e.g. Florisil™).

An "alcohol derivative" is an organic alcohol (ROH, where R is aryl, alkyl or cycloalkyl, and preferably alkyl or cycloalkyl) where the group R may be further substituted with a group selected from, e.g., hydroxy, alkoxy, amino, alkylamino, carboxylic acid, carboxylic ester, keto, aldehyde, nitro, cyano, imino, thio, alkylthio, sulfonic acid or ester, or phosphonic acid or ester. Preferred alcohol derivatives include alcohol-substituted amino acids, glycols, and sugars. Also preferred are hydroxy-terminated polyalkylene oxides, such as polyethylene oxide, having molecular weights in the range of about 300 to 40,000. In general, preferred compounds have at least 5% (w:w) aqueous solubility; however, less soluble alcohols or alcohol derivatives may also be used.

II. Phospholipase-Catalyzed Reactions Employing a Solid Particle Interface

Certain types of solid particles can promote reaction of phospholipids with phospholipases by providing an interface between the lipids, which adsorb to the particles, and the enzyme, which remains predominantly in the aqueous phase. The solid interface may also serve to activate the enzyme. It is possible that immobilization of the enzyme and binding of substrate together on the particle surface results in a change in the conformation of the phospholipase and its conversion to an activated state.

As an example of this method, phospholipase D-catalyzed conversion of egg phosphatidylcholine (PC) to the corresponding phosphatidyl glycerol (PG) was carried out in an aqueous medium using various solid adsorbents in place of the organic solvents (such as diethyl ether) used conventionally. General procedures for small scale reactions are as follows.

Reagents include acetate buffer (pH 5.6) containing 50-200 mM CaCl_2 , glycerol at a concentration of about 25-75% by volume, egg phosphatidylcholine (PC), and phospholipase D. The enzyme may be obtained commercially or prepared as described in Example 1. The substrate (PC) is dispersed in glycerol-acetate buffer, preferably as multilamellar vesicles (MLV), at a concentration of about 5 mg/ml. The MLV may be prepared by hydration of a dry lipid film, the more conventional method, or by reverse evaporation (REV) using a high lipid concentration in a solvent such as ether or a Freon®. In this technique, a nonaqueous solution of vesicle-forming lipids is dispersed with a smaller volume of an aqueous medium to form a water-in-oil emulsion. After removal of the lipid solvent, the resulting gel is converted to liposomes.

Phospholipase D is added to the liposomal dispersion, and the mixture is vortexed until all the enzyme is in solution. The reaction is initiated by adding the solid adsorbent at room temperature, and shaking is commenced immediately.

In one procedure, the reaction is terminated by adding chloroform and methanol to give a final solvent ratio of about 1:1:1 (chloroform:methanol:water). The phases are separated, e.g. by centrifugation, and the phospholipids are isolated from the lower chloroform phase. The products may be separated by preparative TLC chromatography on Unilatch silicic acid glass plates, using a solvent system of chloroform:acetone:methanol:acetic acid:water (6:8:2:2:1).

Alternatively, for preparation of products to be used in food or drugs, where the use of solvents such as chloroform is to be avoided, the lipids may be extracted from the reaction mixture using hexane, heptane, ethanol/hexane or ethanol/heptane mixtures, under either acidic or neutral conditions. If desired, Ca^{+2} ions can be removed from the product lipids by a chelating agent such as EDTA, which will partition into the aqueous phase.

The level of conversion to phosphatidyl glycerol (PG) is determined by TLC followed by phosphorus content analysis, by using a refractive index or light scattering detector, or by the use of radioactively labeled glycerol, as described in Example 2. Amine lipid products may also be assayed colorimetrically by reaction with picryl sulfonate (TNBS).

Results of the reaction using various solid adsorbents, at a ratio of about 70 mg adsorbent per mg lipid, are given in Table 1. The Table shows conversions to PG and to PS in reactions of PC with glycerol and serine, respectively. It is clear that, among the adsorbents tested, silica gel was the most effective.

Table 1 : Activation of Phospholipase D by Solid Adsorbents

Adsorbent	% Conversion (± 5)	
	PG	PS
Silica gel (Kieselgel® 60 Merck)	90	65
Kieselguhr® (BDH)	70	50
Celite® (Koch-light)	40	-
Super-cel® (Amend)	20	-
None	0	0

In further experiments, various types and particle sizes of silica gel and other adsorbents were tested for their activity in the enzymatic conversion of PC to PG. Each reaction system contained 10 mg PC dispersed in 0.5 ml acetate buffer (pH 5.60) containing 25% (vol) glycerol and 50 mM CaCl_2 . Cabbage phospholipase D (0.5 mg, Boehringer) was added, and the reaction was initiated by adding 0.2 grams adsorbent. The mixtures were shaken for 30 minutes at room temperature.

The results are shown in Table 2. As can be seen from this Table, the best conversions were obtained with TLC grade silicic acid (from Merck, Philadelphia, PA, and Camag, Muttenz, Switzerland) (50-60% conversion to PG) and with TLC magnesium silicate from Bio-Rad (Richmond, CA) (50% conversion).

A correlation between the particle size of the silicic acid and conversion was found, with smaller particle sizes giving higher yields. The best results in this series were obtained with silica gel 60H and HR, TLC Grade (Merck), which have the finest particles (particle size distribution 5-20 μ m, mean particle size 10-12 μ m). TLC grade magnesium silicate from Bio-Rad (Richmond, CA) also gave good conversion (particle size distribution 2-44 μ m). A much larger particle size magnesium silicate, used in column chromatography, gave only 20% conversion (Florisil® entry in Table 2). Accordingly, preferred mean particle sizes of silica gel (silicic acid or magnesium silicate) for use in the reactions described herein would be less than 25 μ m, and preferably less than 15 μ m.

With aluminum oxide adsorbents (neutral, acidic or basic alumina) very low conversions were achieved (5-15% PG). When adsorbent materials such as Bio-Beads® SM-2 from Bio-Rad or Chelex® (chelating resin) from Sigma were used, no conversion was obtained.

Table 2: Effect of Different Types and Sizes of Adsorbents on PC \rightarrow PG Conversion

	Adsorbent	PG (\pm 10%)
Silicic Acid	Merck Silica gel H/HR, TLC grade (5-20 μ m)	60
	Merck Silica gel mesh <230 (>60 μ m)	50
	Merck Silica gel mesh 70-230 (60-210 μ m)	40
	Merck Silica gel extra pure mesh 70-230	40
	Merck Silica gel mesh 35-70 (210-500 μ m)	30
	Sigma, Silica gel, mesh 325 (ca. 40 μ m)	40
	Sigma, Silica gel, mesh 60-200 (75-250 μ m)	10
	CAMAG, Silica gel for TLC	50
	Bio-Rad, Bio-Sil® for TLC	40
	BDH, Silica gel, 6-20 mesh (>840 μ m)	5
Mg Silicate	BDH Sand	10
	Bio-Rad (2-44 μ m)	50
Aluminum oxide	Florisil® (column chromatography grade)	20
	Bio-Rad, neutral alumina	15
	Merck, neutral alumina	5
	Sigma, acidic alumina	5
	Basic alumina	0
Bio-Beads®	SM-2 Bio-Rad	0
Chelex®	chelating resin (Sigma)	0

It was also found that, for a given amount of enzyme, the percentage of transphosphatidylolation increased with the amount of silica gel per milligram phospholipid, as shown in Table 3.

Table 3: Effect of Amount of Silica Gel on PC → PG Conversion

Mg adsorbent/mg phospholipid	% PC to PG Conversion ($\pm 10\%$)
20	20
40	60
60	90

The effect of varying the amount of silica gel was further investigated in the 500 mg scale reactions described below. Again, conversion increased with an increase in silica gel:lipid ratio. Conversion appeared to level off at a ratio of approximately 10:1 (Table 4); however, the difference between the last two ratios tested (9/1 and 11/1) is not substantial. Reactions were also performed on a 20g scale at silica/PC ratios of 5 and 25; conversions obtained were 40% and 60%, respectively.

Based on these results, it appears that yields generally increase with silica gel/lipid ratio. Accordingly, ratios of at least 4/1, and more preferably at least 10/1, are preferred. In some cases, such as shown in Table 4, conversion appeared to level off, but no adverse effects of increasing the silica gel/lipid ratio were observed in any case.

Table 4: Effect of Amount of Silica Gel on 500mg Scale PC → PG Conversion

Silica Gel/PC Ratio	PG Formation (% ± 10)
3.3	24
6.7	28
9	40
11	40
Reaction conditions: enzyme concentration = 7.5 mg/ml; buffer = 40 mM acetate, pH 5.6, containing 50% glycerol v/v 100 mM Ca ⁺⁺ ; reaction time = 20 hours, rt.	

II. Effect of Other Reaction Variables on Conversion

The effects of other parameters on the representative egg PC → PG reaction were also studied. These variables included enzyme concentration, glycerol concentration, pH, Ca⁺⁺ concentration, liposome size, order of reagent addition, lipid composition, presence of antioxidants, enzyme source and preparation, and age of enzyme preparation.

Most of the reactions in these studies were carried out on a larger scale, using 500 mg to 5 g of lipid. The following procedure was used for the mid-scale reaction. Egg phosphatidylcholine (500 mg) was dissolved in chloroform, which was then evaporated by a stream of N₂. The dried

lipid was dispersed in 10 ml acetate buffer (40 mM, pH 5.6) containing 50% glycerol (v/v) and CaCl_2 (100 mM). Multilamellar vesicles (MLV) were formed by vigorous shaking of the suspension, using a Lab Line multi wrist shaker (speed setting = 7) at room temperature, until all the lipid was incorporated in the liposomal dispersion. Alternatively, MLV can be prepared by reverse evaporation (REV), as described above.

Phospholipase D (lyophilized powder, 50 mg) was added to the liposomal dispersion and the mixture vortexed until all the enzyme was in solution. The reaction was initiated by adding silicic acid (3 grams, Kieselgel® 60 Merck) at room temperature, and shaking was commenced immediately. The reaction was terminated by adding chloroform and methanol to give a final solvent ratio of 1:1:1 chloroform:methanol:water. Two phases formed after vortexing and were separated by centrifugation at 1000 g. The phospholipids were isolated from the lower chloroform phase by preparative TLC chromatography on Analtech silicic acid glass plates, using a solvent system of chloroform:acetone:methanol:acetic acid:water (6:8:2:2:1).

A. Enzyme Concentration

Table 5 shows the effect of varying enzyme concentration. The optimum concentration under these conditions was about 7 mg/ml. Above this concentration a plateau in the PG formation was obtained.

Table 5. Effect of Enzyme Concentration of PG Formation

Enzyme Concentration (mg/ml)	PG Formation (% \pm 10)
0.5	21
1	31
3	36
7.5	42
15	40
15 (stepwise addition)	40
Reaction conditions: Buffer = 40 mM Acetate, pH 5.6, containing 50% glycerol v/v and 100 mM Ca^{++} ; lecithin/silica gel ratio = 0.15; reaction time = 20 hours; room temperature; reaction volume = 10 ml.	

B. Glycerol Concentration

Effect of glycerol concentration is shown in Table 6. The product, PG, was formed in greater amounts and at a higher rate at lower glycerol concentrations (Table VI). It is possible that high glycerol concentrations increase the viscosity of the incubation medium, thus decreasing the rate of PG formation, or that the glycerol coats the silica gel. However, not unexpectedly, less of the hydrolysis product (PA) was formed at higher glycerol concentrations.

Table 6: Effect of Glycerol Concentration on PG Formation

Glycerol Concentration (% v/v)	PG Formation (% \pm 10)	PA Formation (% \pm 3)
25	70	10
50	40	7
75	15	4
Reaction conditions: enzyme concentration = 7.5 mg/ml; buffer = 40 mM acetate pH 5.6 100 mM Ca^{+2} ; lecithin/silica gel ratio = 0.15; reaction time = 20 hours; room temperature		

5 C. Ca^{+2} Concentration; pH

Based on the data in Table 7, the optimal Ca^{+2} concentration for the phospholipase D-catalyzed transphosphatidyl transfer reaction under these conditions appeared to be about 50-100 mM. However, some preparations worked well at Ca^{+2} concentrations of 5 mM or even lower. Other phospholipases may require different ions, such as Mg^{+2} (for C-type sphingomyelinases) or Zn^{+2} , and some have no ion requirement.

Table 7: Effect of Ca^{+2} Concentration

Incubation Buffer, mM CaCl_2	PG Formation (% \pm 10)	PA Formation (% \pm 3)
50	37	3
100	40	4
150	28	3
200	21	3
Reaction conditions: enzyme concentration = 5 mg/ml lecithin/silica gel ratio = 0.11; buffer = 40 mM Tris pH 8.5; reaction time = 20 hours; room temperature.		

Similar PG conversions were obtained using Tris buffer (pH 8.5) and acetate buffer (pH 5.6).

15 However, somewhat less PA was formed at pH 8.5 (Table 8).

Table 8: Effect of pH

Incubation Buffer	PG Formation (% \pm 10)	PA Formation (% \pm 3)
Acetate, pH 5.6	59	14
Tris, pH 7.5	38	10
Tris, pH 8.5	60	10
Reaction conditions: enzyme concentration = 7.5 mg/ml, lecithin/silica gel ratio = 0.15; reaction time = 20 hours; room temperature.		

20 D. Source of Enzyme

The enzymatic activities of different batches of phospholipase D, prepared in-house from

cabbage as described in Example 1, were compared to the activities of phospholipase D from various commercial sources. Cabbage phospholipase D from Boehringer Co. (Indianapolis, IN) showed a markedly superior activity to the in-house preparations and to phospholipase obtained from Sigma Co. (St. Louis, MO), giving similar yields of PG from PC with only one-fifth of the amount of enzyme (PC/enzyme = 25:1). Phosphatidic acid (PA) was formed with all the enzymes (5-15% relative to total phospholipid).

In view of the superior enzymatic activity of the commercial phospholipase D from the Boehringer Co., additional purification of in-house cabbage phospholipase D preparation was carried out, following the procedure described by Allgyer and Wells. However, no increase in the transphosphatidylase specific activity of the in-house phospholipase was obtained.

E. Age of Enzyme Preparation: Transferase/Hydrolase Activity

It is generally observed that the transferase activity of phospholipase D is less stable than its hydrolase activity. Thus, the ratio of transferase/hydrolase activity of the enzyme tends to diminish with time after its preparation. This variation in transferase/hydrolase activity has been a source of irreproducibility from one preparation of enzyme to another. In reactions conducted by the authors using the biphasic ether/water system, phospholipase D from Savoy cabbage lost its transferase activity by at least 20% per month even upon storage at -20°C, and much more when stored at 4°C in solution.

The transphosphatidylase activity of some old preparations (2-5 years old) of phospholipase D from cabbage, stored at either -70°C or -20°C, was found to be negligible or absent in the above biphasic water/ether system, while the hydrolase activity was largely maintained. However, all of these preparations were able to convert PC to PG in the water/silica gel system. Similar results were obtained with phospholipase D preparations that were stored in aqueous solutions at 4°C. These were totally inactive in the ether/water biphasic system but retained their full activity with silica gel, giving PC to PG conversions of more than 80%.

Because the transferase activity of the enzyme in the conventional ether/water biphasic system decreases with time, the enzyme is typically added to the reaction stepwise. However, in the water/silica gel system, the enzyme appears to be stable throughout the reaction, and may be added either stepwise or all at once with no effect on PG formation.

F. Effect of Direct Adsorption of Lipid onto Adsorbent

The formation of the liposomal suspension can be a time consuming process for a large-scale production. Therefore, the possibility of eliminating this step, by directly adsorbing the lipid from a chloroformic solution onto the silica powder, was examined. For this purpose, two 10 mg samples of PC were dissolved in 4 ml of CHCl₃ in round-bottomed flasks, and two amounts of silica gel 60 H (Merck), 0.1 and 1.0 grams, respectively, were added. The chloroform was evaporated

under reduced pressure. Acetate buffer, glycerol, CaCl_2 and enzyme were added to the dry powder, and the reaction was carried out essentially as described above. The procedure was repeated using magnesium silicate.

TLC chromatograms of samples taken after the reactions were terminated showed no production of PG in any case, indicating that this method of reaction is not viable, and suggesting that the reacting phospholipid must be dispersed in the buffer for the transphosphatidylation to take place.

G. Other Factors: Activation/Deactivation of Adsorbents; Liposome Size; Antioxidant

These factors were found to have little or no effect on conversion in the above-described reaction. To test the effect of adsorbent activation/deactivation, the two types of adsorbents giving the best results in the reaction, i.e. silicic acid 60 H (Merck) and magnesium silicate (Florisil®) (Bio-Rad), were submitted to heat activation (dehydration) and deactivation (hydration). The activation was performed by heating the adsorbents at 100°C overnight and the deactivation by adding 0.2 ml of water homogeneously to 200 mg of the adsorbent powder. Conversions obtained with these treated adsorbents were compared with non-treated adsorbents (Example 3). TLC chromatograms showed similar conversions in all the systems tested, in the range of 50-60% PG production. It can therefore be concluded that dehydration or hydration of the adsorbent does not affect conversion, so there is no need for pretreatment of the adsorbent before the reaction.

Based on the assumption that conversion may be limited by the exposure of substrate to enzyme, the influence of the liposome size on the efficiency of the transphosphatidylation reaction was examined, using MLV (approx. $1.5\ \mu\text{m}$ diameter) and SUV ($20\text{-}40\ \text{nm}$ diameter). The liposomal dispersions were formed and the reactions carried out as described in Example 4. About 40% PG production was observed in both systems, indicating that substrate availability is not limited by vesicle size.

The peroxidation of phospholipids having polyunsaturated acyl chains can be inhibited by the use of antioxidants. It was found that presence of the antioxidant butylated hydroxytoluene (BHT) did not affect the activity of silica-gel activated phospholipase D in the transesterification reaction. The antioxidant afforded excellent protection against lipid peroxidation during the reaction, allowing the production of high quality PG.

III. Further Scaleup

Reaction at a 5 gram scale (using egg PC prepared from fresh eggs) gave similar results to those at 500 mg scale. The scale was then increased to 20 grams, using commercial phospholipid from the Asahi Co. (Japan). Using the preferred conditions described in Section II above, but reducing the amount of silica gel to a silica gel/lecithin ratio of 4.0, a conversion of approximately 70% PC to PG was obtained after 4 hours of reaction. Most of the conversion occurred during the

first hour. The level of PA was very low ($< 5\%$).

The reaction was also tested under industrial scale conditions using pilot plant reactors. A 100 gram scale process was carried out at the minipilot plant unit of the Casali Institute of Applied Chemistry, School of Applied Sciences, The Hebrew University, Jerusalem, as described in

- 5 Example 5. The PG conversion was 40% at room temperature and 50% at 35°C. About 7% of PA was produced at room temperature, while at 35°C a higher level of PA (15%) was observed.

IV. Variation of Lipid Substrate

- Saturated phospholipids may be more suitable than unsaturated phospholipids for certain drug
10 delivery systems, due to their higher resistance to lipid peroxidation. Partially hydrogenated (PH) egg PC is a potential candidate to replace egg PC. Therefore, the transphosphatidylation of PC to PG was tested with a saturated lipid, dipalmitoyl PC (DPPC; Avanti Polar Lipids, Birmingham, AL) and with partially hydrogenated (PH) PC (iodine value 30; Asahi Chemical, Japan). Reactions
15 were carried out as described in Example 6. TLC of the DPPC reaction showed 50% conversion to DPPG and no trace of PA. A yield of 77% PH-egg PG was obtained from PH-PC, with only traces of PA ($< 5\%$) detected.

- The results of these reactions and of additional substrates and hydroxyl-containing reagents are summarized in Table 9. The reaction gives good conversion for a variety of reagents and substrates, including tissue extracts. Phosphatidyl serine enrichment of a brain phospholipid
20 extract (sixth row of Table 9) is described in more detail in Example 7.

Table 9

Lipid Substrate	Reagent	Product	Typical Conversion
Egg lecithin (PC)	Glycerol	PG	50-70%
Egg lecithin	Serine	PS	70-90%
Egg lecithin	Water	PA	70-100%
Egg lecithin	Myo-inositol	PI	60-80%
Phosphatidyl ethanolamine	Glycerol	PG	15-40%
Brain phospholipid extract	Serine	PS enrichment	80%
Dioleoyl PC	Glycerol	Dioleoyl PG	50-70%
Dipalmitoyl PC	Glycerol	Dipalmitoyl PG	50%
Partially hydrogenated egg PC	Glycerol	PH-PG	75-80%

25 V. Reaction of Other Phospholipases

The biphasic water/silica gel method was employed for reaction of other phospholipases (C and A2) in selective hydrolysis of egg PC. Reactions were done on a 10 g scale, using Merck silica

gel, 70-230 mesh. Other conditions and conversions (after 4 hrs at room temperature) are given in Table 10. Reaction with phospholipase C, under similar conditions, was also used to prepare ceramide from sphingomyelin (data not shown).

Table 10

Phospholipase	Source of enzyme	Medium	Product	Typical Conversion
C	Clostridium welchii (Sigma)	Tris buffer (40mM, pH 7.6) 20 mM Ca^{+2}	Diacyl glycerol	100%
C	Bacillus cereus (Makor)	"	"	70-90%
A2	Snake venom Naja-Naja (Sigma)	Acetate buffer (40mM, pH 5.6) 100 mM Ca^{+2}	Lyso-lecithin	100%
A2	Crotalus adamaneus (Sigma)	"	"	100%

EXAMPLES

The following examples illustrate but are not intended in any way to limit the invention.

Example 1. Isolation of Phospholipase D from Cabbage

A. Standard Preparation

Phospholipase D was prepared from Savoy cabbage according to the procedure of Yang.

A crude extract was first prepared by homogenizing 4250 kg of light-green leaves of fresh Savoy cabbage with distilled water in a Waring blender. The fibrous material was removed by filtering through a gauze. The final volume of the cabbage juice (3750 liters) was then centrifuged at 500 g for 5 minutes at 4°C. The extract was heated to 55°C, maintained at this temperature for 5 minutes, and then rapidly cooled. The bulky precipitate was removed by centrifugation under the same conditions and discarded.

Two volumes of acetone were added to the supernatant at -15°C, and the mixture was shaken immediately and stored overnight at 4°C. The acetone was evaporated and the aqueous suspension was centrifuged. The supernatant was discarded and the precipitate lyophilized overnight under 150 millitorr vacuum. The enzyme, obtained as a light green powder (5.2 g), was stored frozen at -70°C until needed.

It was later found that the yield of enzyme powder could be improved by reducing the volume of water used for the homogenization of the cabbage leaves. The modified procedure yielded 2.6 grams of lyophilized protein per kilogram of cabbage.

B. Simplified Preparation

It was found that by replacing the slow speed centrifugation step (500 g × 15 min. at 4°C) of the above method with high speed centrifugation (15000 g × 15 min. at 4°C), the supernatant could be used as the enzyme preparation without further purification. The high speed centrifugation yields reproducible enzyme preparations free of particulate and fibrous matter. No cabbage residual lipids were detected in the extract, using the method of Bligh and Dyer. The centrifugation step could be further optimized to fit industrial needs by using continuous flow centrifugation. Using this crude fresh enzyme, similar conversions of PC to PG were obtained at similar equivalents of protein weight per cabbage weight using the experimental conditions described above.

Example 2. Determination of Conversion

Lipids are extracted from the incubation mixture by vortexing with a 1:1:1 mixture of chloroform:methanol:water and allowing the phases to separate. The upper aqueous-methanolic phase contains all the water soluble reagents, while the lower chloroformic phase contains the lipids (e.g. PC, PG and phosphatidic acid (PA)). Aliquots of the lower phase are loaded on Analtech silicic acid thin layer on a glass plate for TLC. The phospholipids are separated (Fig. 1) using a solvent system of chloroform:acetone:methanol:acetic acid:water (6:8:2:2:1 v/v). The individual spots are scraped into phosphorous free test tubes, and the phosphorous content is determined using the perchloric acid based Bartlett procedure (see e.g. Barenholz *et al.*, 1993).

An alternative assay for assessing conversion, based on the use of a radioactive water soluble reagent, was developed during the course of this work. For example, tritiated glycerol or serine is used; i.e. ^3H -glycerol (or serine) + PC \rightarrow ^3H -PG (or ^3H -PS) + choline (+PA). The glycerol or serine partitions into the upper methanolic/aqueous phase in the biphasic solvent system of chloroform:methanol: water (8:4:3 or 1:1:1) during workup; all lipids partition into the chloroform lower phase. ^3H -PG or ^3H -PS is determined by scintillation counting of the chloroform phase.

This assay was tested under various conditions and proved to be simpler and much faster than thin layer chromatography. However, it does not determine the level of PA (undesired product).

A spectrophotometric method suitable for the determination of amine phospholipids (i.e. PS or PE) utilizes the reaction of the amine phospholipid with trinitro benzene sulfonate (TNBS) to form the yellow trinitrophenyl derivative (i.e. TNP-PS). This product is determined spectrophotometrically in the chloroform enriched lower phase (Barenholz *et al.*, 1993) or by TLC (Amselem *et al.*, 1993).

Example 3. Conversion of PC to PG using Activated or Deactivated Adsorbent

Egg PC (10 mg) was dispersed in 0.5 ml of 50 mM acetate buffer (pH 5.6) containing 50 mM

CaCl₂ and 50% (vol) glycerol. Cabbage phospholipase D (Boehringer) was added (0.5 mg), and the reaction was started by adding 200 mg of silicic acid or Mg-silicate as normal powder, activated or deactivated. The reaction mixtures were shaken at room temperature for 30 minutes. The reactions were terminated by adding 0.5 ml of distilled water and 2 ml of a CHCl₃:methanol mixture (1:1). The lipids were extracted from the lower phase after centrifugation.

Example 4. Reaction of PC in SUV and MLV

The reactions were carried out on a 5 gram (PC) scale. The lipid was dissolved in 100 ml CHCl₃. Glass beads (100 grams) of 5 mm diameter were added to increase the surface area of the dried lipid film and to ensure a better dispersion of the lipid in the aqueous solution. The organic solvent was removed by a flash evaporator. Then 50 ml of the acetate buffer containing 50% glycerol was added, and the MLV were formed by shaking the mixture in a Lap-line multi-wrist shaker until an homogeneous liposomal dispersion was obtained. The SUV were prepared from these MLV by sonicating the MLV using a probe sonicator (350 heat systems, Ultrasonics Inc.) for 5 minutes. At the same time, 600 grams of cabbage were homogenized in a Waring blender with 250 ml acetate buffer, filtered through a gauze and centrifuged for 20 minutes at 4°C at 15,000 rpm. Fifty mL of this fresh crude supernatant were mixed with 25 grams glycerol and added to the MLV and to the SUV. The reactions were initiated by adding 25 grams silica gel 60 H, the mixture was shaken overnight at room temperature in the multi-wrist shaker.

Example 5. Large Scale Conversion of PC to PG

MLV were prepared by the thin lipid film procedure. Egg phosphatidylcholine (100 grams) from Asahi Co. (Japan) was dissolved in 500 ml CHCl₃ in a 1 liter round bottomed flask. The antioxidant BHT (butylated hydroxytoluene) was added at a molar ratio of 1/1000. Glass beads (400 gram) were added, and the organic solvent was evaporated until dryness. Acetate buffer (pH 5.6, 1 liter) containing 50% glycerol and 0.1 M CaCl₂ was added, and MLV were formed by vigorous shaking of the mixture for 1 hour with the aid of the multi-wrist shaker. The final volume of the MLV prepared was divided into two 0.5 liter portions. To each portion was added 250 ml additional acetate buffer, and the mixtures were introduced in two minireactors of 1.5 liter capacity. Fresh crude cabbage phospholipase D juice (250 ml), diluted 1:1 with glycerol to give a final concentration of 50% glycerol, was added to each minireactor. The temperatures in the two reactors were room temperature (20°C) and 35 ± 5°C, respectively. Silica gel 60 H (Merck) (250 g) was added to each reactor. Mixing of reagents in the reactors with mechanical shaking. The reactions were carried out under a nitrogen atmosphere. After 20 hours of reaction, the shaking was stopped, and 1 ml samples were taken from the bottom of each reactor. Each sample was extracted by adding 1 ml DDW and 2 ml of 1:1 CHCl₃:methanol. The phases were separated by

centrifugation and aliquots of the lower phases were analyzed by TLC, eluting with CHCl_3 :acetone:acetic acid: H_2O (6:8:2:2:1). The spots were scraped and extracted and the phospholipid content determined by the Bartlett procedure.

5 Example 6. Reaction of Saturated and Partially Hydrogenated Lipids

MLV were prepared, using 10 mg DPPC in 10 ml acetate buffer (pH 5.6) containing 50% glycerol and 100 mM CaCl_2 at 55°C (10°C above T_m). Ten mL phospholipase D solution, prepared in house, were mixed with 5 grams glycerol and added to the DPPC-MLV. The reaction was started by adding 2.5 grams silica gel 60 H, and the mixture was shaken for one hour at room temperature.

10 PH-egg PC lipid (5 grams) was dried from chloroform solution on 100 gram glass beads (5 mm diameter). Acetate buffer (450 ml, containing glycerol and CaCl_2 as described above) was added, and MLV were formed by shaking the mixture vigorously for 1 hour with a Lab-line multi-wrist shaker. A solution of 20 mg of cabbage phospholipase D (Boehringer) dissolved in 5 ml
15 acetate buffer was then added. The reaction was started by adding 125 grams silica gel 60 H (Merck), the mixture was shaken overnight at room temperature.

Example 7: Phosphatidyl Serine Enrichment of Brain Phospholipid Extract

Bovine brain phospholipid extract containing 15 mole% PS, 43 mole% PC, and 34 mole% PE
20 was used. MLV were prepared from the extract as described in example 5. L-Serine (powder) was added to the liposomal dispersion at a level close to saturation. Fresh crude cabbage juice (Example 1 simplified procedure) was used as the enzyme source. The reaction was carried out as described in Example 5. Analysis by the TNBS method and by the TLC method, described above, showed that the level of PS was increased from 15 mole% in the starting material to 51 mole% in
25 the final product.

A similar reaction using D,L-serine gave 49 mole% phosphatidyl serine in the final product.

While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications may be made without departing from the invention.

IT IS CLAIMED:

1. A method of conducting an enzyme-catalyzed transesterification or hydrolysis of a phospholipid, comprising

- 5 dissolving said enzyme in an aqueous medium containing (i) a liposomal suspension of said phospholipid, a (ii) hydroxyl-containing reagent selected from water, an alcohol or an alcohol derivative, and (iii), when required by the enzyme, a divalent metal cation, adding silica gel to said medium, and agitating the resulting mixture.

- 10 2. The method of claim 1, wherein the enzyme is selected from the group consisting of phospholipase A1, phospholipase A2, phospholipase C, phospholipase D, and a sphingomyelin phosphodiesterase.

- 15 3. The method of claim 2, wherein the enzyme is selected from the group consisting of phospholipase A2, phospholipase C, and phospholipase D.

4. The method of claim 3, wherein the enzyme is phospholipase D.

- 20 5. The method of claim 4, wherein the divalent metal cation is calcium ion in concentration of about 5-100 mM.

6. The method of claim 1, wherein the hydroxyl-containing reagent is an alcohol or alcohol derivative.

- 25 7. The method of claim 6, wherein the hydroxyl-containing reagent is selected from the group consisting of glycerol, serine, and inositol.

8. The method of claim 6, wherein the hydroxyl-containing reagent is a hydroxy-terminated polyethylene glycol having a molecular weight between about 300 and 40,000.

30

9. The method of claim 1, wherein the silica gel is added in an amount which is at least four times the amount of the phospholipid by weight.

10. The method of claim 9, wherein the silica gel is added in an amount which is at least ten times the amount of the phospholipid by weight.

35

11. The method of claim 1, wherein the silica gel has a mean particle size no greater than

WO 00/77183

PCT/IL00/00350

25 μ m.

12. The method of claim 11, wherein the silica gel has a mean particle size no greater than 15 μ m.

5

13. The method of claim 1, wherein the phospholipase is present in said medium at a concentration of at least 3 mg/ml.

14. The method of claim 13, wherein the phospholipase is present in said medium at a concentration of at least 7 mg/ml.

10

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 December 2000 (21.12.2000)

PCT

(10) International Publication Number
WO 00/77183 A1

(51) International Patent Classification: C12N 9/16,
C12P 9/00, 13/00, C12N 9/18

(21) International Application Number: PCT/IL00/00350

(22) International Filing Date: 15 June 2000 (15.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/139,316 15 June 1999 (15.06.1999) US

(71) Applicant (for all designated States except US): YISSUM
RESEARCH DEVELOPMENT COMPANY OF THE
HEBREW UNIVERSITY OF JERUSALEM [IL/IL];
Jabotinsky Street 46, 91042 Jerusalem (IL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BARENHOLZ,
Yechezkel [IL/IL]; Nave Shanan Street 18, 93707
Jerusalem (IL). AMSELEM, Shimon [IL/IL]; Miriam
Mizrachi Street 10, 76551 Rehovot (IL).

(74) Agent: REINHOLD COHN AND PARTNERS; P.O.
Box 4060, 61040 Tel-Aviv (IL).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guidance
Notes on Codes and Abbreviations" appearing at the beginning
of each regular issue of the PCT Gazette.

(54) Title: ENZYMATIC PREPARATION OF PHOSPHOLIPIDS IN AQUEOUS MEDIA

(57) Abstract: Methods of conducting phospholipase-catalyzed transesterification or hydrolysis of phospholipids present in aqueous liposomal suspensions, in the absence of detergents and organic solvents, are described. The method, which employs a water/solid particle interface, gives high conversions, particularly when the solid particle is silica gel having a small particle size and is present at a level at least four times the weight of the reacting lipid. The reaction is useful for the preparation of a variety of differently substituted phospholipids, as well as diacyl glycerols and ceramides.

WO 00/77183 A1



Combined Declaration for Patent Application and Power of Attorney

I, a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

ENZYMATIC PREPARATION OF PHOSPHOLIPIDS IN AQUEOUS MEDIA

the specification of which (check one)

☐ is attached hereto;

☐ was filed in the United States under 35 U.S.C. §111 on, as
U.S. Appl. No. _____, *

[X] was filed in the U.S. under 35 U.S.C. §371 by entry into the U.S. national stage of an international (PCT) application, PCT/IL00/00350; filed June 15, 2000, entry requested on December 17, 2001*; national stage application received U.S. Appl. No. 10/009,771, *, §371/§102(e) date _____ * (* if known)

and was amended on December 17, 2001 (if applicable).

(include dates of amendments under PCT Art. 19 and 34 if PCT)

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and I acknowledge the duty to disclose to the Patent and Trademark Office (PTO) all information known by me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119 (a)-(d) and 365 (b) of any prior foreign application(s) for patent, inventor's or plant breeder's rights certificate(s), or under §365(a) of any PCT application which designated at least one country other than the U.S., listed below:

Application No.	Country	Filing Date (MM/DD/YYYY)
-----------------	---------	--------------------------

If I claimed foreign priority above, I hereby identify below any foreign application for patent (including an international (PCT) application designating a country other than the United States) or for an inventor's or plant breeder's certificate, having a filing date before that of the earliest application from which foreign priority is claimed (if left blank, then there are none):

Non-Priority Application No.	Country	Filing Date (MM/DD/YYYY)
------------------------------	---------	--------------------------

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional applications listed below:

Application No.	Filing Date (MM/DD/YYYY)
60/139,316	June 15, 1999

I hereby claim the benefit under 35 U.S.C. §120 of any prior U.S. non-provisional application(s) or under §365(i) of any prior PCT international application(s) designating the U.S., listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in such U.S. or PCT international application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the PTO all information which is material to patentability as defined in 37 C.F.R. §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Application No.	Filing Date (MM/DD/YYYY)	Status (patented, pending, abandoned)
PCT/H00/00350	June 15, 2000	Pending

As a named inventor, I hereby appoint the following registered practitioners to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

All of the practitioners associated with Customer Number 001444

Direct all correspondence to the address associated with Customer Number 001444, which is presently:

BROWDY AND NEIMARK, P.L.L.C.
624 Ninth Street, N.W.
Washington, D.C. 20001-5303
(202) 628-5197

Title: ENZYMATIC PREPARATION OF PHOSPHOLIPIDS IN AQUEOUS MEDIAU.S. Application filed December 17, 2001, Serial No. _____PCT Application filed June 15, 2000, Serial No. PCT/IL00/00350

The undersigned hereby authorizes the U.S. Attorneys or Agents appointed herein to accept and follow instructions from Reinhold Cohn & Partners as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorneys or Agents and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents appointed herein will be so notified by the undersigned.

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1-00 FULL NAME OF FIRST INVENTOR <u>Yechezkel BARENHOLZ</u>		INVENTOR'S SIGNATURE <u>Yechezkel Barenholz</u> <u>ILX</u>	DATE <u>8.12.2007</u>
RESIDENCE <u>Jerusalem, Israel</u>		CITIZENSHIP <u>Israel</u>	
POST OFFICE ADDRESS <u>Nave Shanan Street 18, 93707 Jerusalem, Israel</u>			
FULL NAME OF SECOND JOINT INVENTOR <u>Shimon AMSELEM</u>		INVENTOR'S SIGNATURE	DATE
RESIDENCE <u>Rehovot, Israel</u>		CITIZENSHIP <u>Israel</u>	
POST OFFICE ADDRESS <u>Mizrachi Street 10, 76551 Rehovot, Israel</u>			
FULL NAME OF THIRD JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF FOURTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF FIFTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF SIXTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP	
POST OFFICE ADDRESS			

ALL INVENTORS MUST REVIEW APPLICATION AND DECLARATION BEFORE SIGNING. ALL ALTERATIONS MUST BE INITIALED AND DATED BY ALL INVENTORS PRIOR TO EXECUTION. NO ALTERATIONS CAN BE MADE AFTER THE DECLARATION IS SIGNED. ALL PAGES OF DECLARATION MUST BE SEEN BY ALL INVENTORS.

Page 2 of 2 Pages

Atty. Docket: BARENHOLZ-6

Title: ENZYMATIC PREPARATION OF PHOSPHOLIPIDS IN AQUEOUS MEDIAU.S. Application filed December 17, 2001

Serial No.

PCT Application filed June 15, 2000Serial No. PCT/IL00/00350

The undersigned hereby authorizes the U.S. Attorneys or Agents appointed herein to accept and follow instructions from Reinhold Colm & Partners as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorneys or Agents and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents appointed herein will be so notified by the undersigned.

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, did that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1-00	FULL NAME OF FIRST INVENTOR <u>Yecheskel BARENHOLZ</u>	INVENTOR'S SIGNATURE <u>Yecheskel Barenholz</u>	DATE <u>12.02.02</u>
	RESIDENCE <u>Jerusalem, Israel</u>	CITIZENSHIP	
	POST OFFICE ADDRESS <u>Nave Shanan Street 18, 93707 Jerusalem, Israel</u>		
2-00	FULL NAME OF SECOND JOINT INVENTOR <u>Shimon AMSELEM</u>	INVENTOR'S SIGNATURE <u>Shimon Amselem</u>	DATE <u>19.2.02</u>
	RESIDENCE <u>Rehovot, Israel</u>	CITIZENSHIP <u>Israel</u>	
	POST OFFICE ADDRESS <u>Mizrachi Street 10, 76551 Rehovot, Israel</u>		
	FULL NAME OF THIRD JOINT INVENTOR	INVENTOR'S SIGNATURE	DATE
	RESIDENCE	CITIZENSHIP	
	POST OFFICE ADDRESS		
	FULL NAME OF FOURTH JOINT INVENTOR	INVENTOR'S SIGNATURE	DATE
	RESIDENCE	CITIZENSHIP	
	POST OFFICE ADDRESS		
	FULL NAME OF FIFTH JOINT INVENTOR	INVENTOR'S SIGNATURE	DATE
	RESIDENCE	CITIZENSHIP	
	POST OFFICE ADDRESS		
	FULL NAME OF SIXTH JOINT INVENTOR	INVENTOR'S SIGNATURE	DATE
	RESIDENCE	CITIZENSHIP	
	POST OFFICE ADDRESS		

ALL INVENTORS MUST REVIEW APPLICATION AND DECLARATION BEFORE SIGNING. ALL ALTERATIONS MUST BE INITIALED AND DATED BY ALL INVENTORS PRIOR TO EXECUTION. NO ALTERATIONS CAN BE MADE AFTER THE DECLARATION IS SIGNED. ALL PAGES OF DECLARATION MUST BE SIGNED BY ALL INVENTORS.

RECEIVED

MAY - 8 2002

TECHNOLOGY CENTER R3700

UNISSIA